

## Sesbania Aculeata, Danch - 0.5 Kg Seeds



It is known by many common names, including danchi, dunchi, dhaincha, canicha, prickly sesban, ""Jantar"" or spiny sesbania

Rating: Not Rated Yet

**Price**

Sales price R 1,607

Discount

[Ask a question about this product](#)

Description The legume *Sesbania bispinosa*, also known as *Sesbania aculeata* Pers., is a small tree in the genus *Sesbania*. The flower is the provincial flower of Phra Nakhon Si Ayutthaya Province, Thailand.

**Common name:** *Sesbania aculeata*, *Aeschynomene aculeata*

**Color:** green

**Bloom time:** june to august

**Difficulty level:** easy to grow

**Planting & Care**

Plant where they will receive a minimum of 5 to 6 hours of full sun per day. Plants grown in weak sun may not die at once, but they weaken gradually. Give them plenty of organic matter when planting and don't crowd them. Wear sturdy gloves to protect your hands from prickly thorns.

**Sunlight:** full sun

**Soil:** well-drained soil

**Water:** Medium

**Temperature:** 75 C

**Fertilizer:** Apply any organic fertilizer

**Care:**

- Enrich the soil with leaf mold or well-rotted organic matter to encourage good drainage.
- Dig a hole 2 to 3 times as deep as the bulbs are high and set the bulb in the hole pointy side up.
- Fill the hole with soil and tamp gently.
- Space bulbs at a distance equal to 3 times the bulb's diameter.
- For a good effect, plant plants in groups of 3 to 5 bulbs.

**Harvesting:** 6-8 days

**Care:**

- Feed plants on a regular basis before and throughout the blooming cycle (avoid chemical fertilizers and pesticides if you're harvesting for the kitchen).
- Once a month between April and July, apply a balanced granular fertilizer (5-10-5 or 5-10-10).
- Allow  $\frac{3}{4}$  to 1 cup for each bush, and sprinkle it around the drip line, not against the stem.

**Special Feature:**

Green manuring of rice with dhaincha (*Sesbania aculeata*) is widely practised under irrigated puddle-transplanted conditions. In flood-prone lowlands, the rice is established through direct seeding early in the season and flooding occurs after 1–2 months of crop growth following regular rains.

Use

**Note:**

for medicinal use

**Reference:**

<http://www.missouribotanicalgarden.org/> <http://www.flowersofindia.net/>

**Reviews**

There are yet no reviews for this product.